

MSLN/Mesothelin
Catalog # PVGS1749**Specification**

MSLN/Mesothelin - Product Information

Primary Accession [Q61468-1](#)
Species
Mouse

Sequence
Asp298-Ser600

Purity
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC

Endotoxin Level
Less than 1EU per µg by the LAL method.

Biological Activity
Immobilized MSLN/Mesothelin(298-600), His, Mouse(Cat.No.: Z03876) at 2 µg/ml(100 µl/Well) on the plate can bind Biotinylated Human CA125, His Tag.

Expression System
HEK293

Theoretical Molecular Weight
35.2 kDa

Formulation **Lyophilized from a 0.22 µm filtered solution in PBS, pH 7.4.**

Reconstitution
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.

Storage & Stability
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

MSLN/Mesothelin - Additional Information

Target Background
Mesothelin, also known as MSLN, is encoded by the MSLN gene. It encodes a precursor protein of 71 kDa that is processed to a 31 kDa shed protein called megakaryocyte potentiating factor (MPF) and a 40 kDa fragment, mesothelin. MSLN is attached to the cell membrane by a glycosyl-phosphatidylinositol (GPI) anchor. It is a differentiation antigen which is highly expressed in several human cancers, including virtually all mesotheliomas and pancreatic adenocarcinomas.

MSLN/Mesothelin - Protein Information

MSLN/Mesothelin - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MSLN/Mesothelin - Images