

**DLL4**  
**Catalog # PVGS1825****Specification**

---

**DLL4 - Product Information**

Primary Accession [Q9NR61](#)  
**Species**  
Human

**Sequence**  
Ser27-Pro524

**Purity**  
> 95% as determined by Bis-Tris PAGE  
> 95% as determined by HPLC

**Endotoxin Level**  
Less than 1EU per µg by the LAL method.

**Biological Activity**  
Immobilized DLL4[Biotin], His, Human (Cat.No.: Z03956) at 2 µg/ml (100 µl/Well) on the streptavidin precoated plate can bind Anti-DLL4 Antibody, hFc Tag

**Expression System**  
HEK293

**Theoretical Molecular Weight**  
55.7 kDa

Formulation **Lyophilized from a 0.22 µm filtered solution in PBS, 200mM L-arginine(pH 7.4).**

**Reconstitution**  
It is recommended that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute the lyophilized powder in ddH<sub>2</sub>O more than 100 µg/ml.

**Storage & Stability**  
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.

**DLL4 - Additional Information**

**Gene ID** 54567

**Other Names**  
Delta-like protein 4, Drosophila Delta homolog 4, Delta4, DLL4

**Target Background**  
Delta-like protein 4 (DLL4) is a type I transmembrane protein with a DSL domain and eight tandem EGF repeats. DLL4 functions as a Notch ligand and activates NOTCH1 and NOTCH4 in the Notch

signaling pathway. It is involved in vascular development and homeostasis. DLL4 is involved in vascular development and homeostasis. It is highly expressed in some cancers, such as bladder, breast cancers.

## **DLL4 - Protein Information**

### **Name** DLL4

### **Function**

Involved in the Notch signaling pathway as Notch ligand (PubMed:<a href="http://www.uniprot.org/citations/11134954" target="\_blank">11134954</a>). Activates NOTCH1 and NOTCH4. Involved in angiogenesis; negatively regulates endothelial cell proliferation and migration and angiogenic sprouting (PubMed:<a href="http://www.uniprot.org/citations/20616313" target="\_blank">20616313</a>). Essential for retinal progenitor proliferation. Required for suppressing rod fates in late retinal progenitors as well as for proper generation of other retinal cell types (By similarity). During spinal cord neurogenesis, inhibits V2a interneuron fate (PubMed:<a href="http://www.uniprot.org/citations/17728344" target="\_blank">17728344</a>).

### **Cellular Location**

Cell membrane; Single-pass type I membrane protein

### **Tissue Location**

Expressed in vascular endothelium.

## **DLL4 - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## **DLL4 - Images**