

CD7
Catalog # PVGS1863**Specification**

CD7 - Product InformationPrimary Accession [XP_005585387.1](#)**Species**
Cynomolgus**Sequence**
Ala26-Pro180**Purity**
> 95% as determined by Bis-Tris PAGE
> 95% as determined by HPLC**Endotoxin Level**
Less than 1EU per µg by the LAL method.**Biological Activity**
Measured by its binding ability in a functional ELISA. Immobilized CD7, His, Cynomolgus at 1 µg/ml (100 µl/well) on the plate can bind Human SECTM1, hFc Tag. Test result was comparable to standard batch.**Expression System**
HEK293**Theoretical Molecular Weight**
17.43 kDa**Formulation**
Lyophilized from a 0.22 µm filtered solution in PBS (pH 7.4).**Reconstitution**
Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.**Storage & Stability**
Upon receiving, the product remains stable up to 6 months at -20 °C or below. Upon reconstitution, the product should be stable for 3 months at -80 °C. Avoid repeated freeze-thaw cycles.**CD7 - Additional Information****Target Background**
CD7, also known as Leu-9, is an approximately 40 kDa glycosylated and palmitoylated transmembrane protein in the immunoglobulin superfamily. CD7 is expressed on T cells, NK cells, myeloid progenitor cells, and CD19 B progenitor cells. Among CD8 T cells, the CD7-bright population preferentially contains naïve and memory cells, while more weak expressors are primarily effector cells.

CD7 - Protein Information

CD7 - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD7 - Images