

SARS-CoV-2 Spike protein (S1)
Catalog # PVGS1955**Specification**

SARS-CoV-2 Spike protein (S1) - Product InformationPrimary Accession [P0DTC2](#)**Species**
SARS-CoV-2**Sequence**
Gln14-Arg685**Purity**
> 90% as analyzed by SDS-PAGE**Endotoxin Level**
< 0.2 EU/ µg of protein by gel clotting method**Biological Activity**
SARS-CoV-2 Spike protein (S1) can bind with Human ACE2 in functional ELISA assay.**Expression System**
293 Cells**Theoretical Molecular Weight**
78.3 kDa
(Additional amino acids from the vector are included. The sequence of them is confidential.)Formulation **Supplied as a solution in PBS pH 7.4****Storage & Stability**

Upon receiving, this product remains stable for up to 6 months at -20°C or below. Avoid repeated freeze-thaw cycles.

SARS-CoV-2 Spike protein (S1) - Additional Information**Gene ID** 43740568**Other Names**

Spike glycoprotein {ECO:0000255|HAMAP-Rule:MF_04099}, S glycoprotein {ECO:0000255|HAMAP-Rule:MF_04099}, E2 {ECO:0000255|HAMAP-Rule:MF_04099}, Peplomer protein {ECO:0000255|HAMAP-Rule:MF_04099}, Spike protein S1 {ECO:0000255|HAMAP-Rule:MF_04099}, Spike protein S2 {ECO:0000255|HAMAP-Rule:MF_04099}, Spike protein S2' {ECO:0000255|HAMAP-Rule:MF_04099}, S {ECO:0000255|HAMAP-Rule:MF_04099}

Target Background

SARS-CoV-2 (Severe Acute Respiratory Syndrome Coronavirus 2) also known as 2019-nCoV (2019 Novel Coronavirus) is a virus that causes illnesses ranging from the common cold to severe diseases. SARS-CoV-2 Spike Protein is composed of S1 domain and S2 domain. S1 contains a

receptor-binding domain (RBD) that can specifically bind to angiotensin-converting enzyme 2 (ACE2), the receptor on target cells. S protein plays an important role in the induction of neutralizing-antibodies and T-cell responses, as well as protective immunity.

SARS-CoV-2 Spike protein (S1) - Protein Information

Name S {ECO:0000255|HAMAP-Rule:MF_04099}

Function

[Spike protein S1]: Attaches the virion to the cell membrane by interacting with host receptor, initiating the infection. The major receptor is host ACE2 (PubMed:32142651, PubMed:32155444, PubMed:33607086). When S2/S2' has been cleaved, binding to the receptor triggers direct fusion at the cell membrane (PubMed:34561887). When S2/S2' has not been cleaved, binding to the receptor results in internalization of the virus by endocytosis using host TFRC and GRM2 and leading to fusion of the virion membrane with the host endosomal membrane (PubMed:32075877, PubMed:32221306, PubMed:34903715, PubMed:36779763). Alternatively, may use NRP1/NRP2 (PubMed:33082294, PubMed:33082293) and integrin as entry receptors (PubMed:35150743). The use of NRP1/NRP2 receptors may explain the tropism of the virus in human olfactory epithelial cells, which express these molecules at high levels but ACE2 at low levels (PubMed:33082293). The stalk domain of S contains three hinges, giving the head unexpected orientational freedom (PubMed:32817270).

Cellular Location

Virion membrane {ECO:0000255|HAMAP-Rule:MF_04099, ECO:0000269|PubMed:32979942}; Single-pass type I membrane protein {ECO:0000255|HAMAP-Rule:MF_04099, ECO:0000269|PubMed:34504087}. Host endoplasmic reticulum-Golgi intermediate compartment membrane {ECO:0000255|HAMAP-Rule:MF_04099, ECO:0000269|PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255|HAMAP-Rule:MF_04099}. Host cell membrane {ECO:0000255|HAMAP-Rule:MF_04099, ECO:0000269|PubMed:34504087}; Single-pass type I membrane protein {ECO:0000255|HAMAP-Rule:MF_04099}. Note=Accumulates in the endoplasmic reticulum-Golgi intermediate compartment, where it participates in virus particle assembly. Some S oligomers are transported to the host plasma membrane, where they may mediate cell-cell fusion (PubMed:34504087). An average of 26 +/-15 S trimers are found randomly distributed at the surface of the virion (PubMed:32979942) {ECO:0000255|HAMAP-Rule:MF_04099, ECO:0000269|PubMed:32979942, ECO:0000269|PubMed:34504087}

SARS-CoV-2 Spike protein (S1) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SARS-CoV-2 Spike protein (S1) - Images