

**Biotinylated p21Cip1-T145 Phospho Peptide**  
**Synthetic Peptide**  
**Catalog # SP2035b****Specification**

---

**Biotinylated p21Cip1-T145 Phospho Peptide - Product Information**

Primary Accession	<a href="#">P38936</a>
Other Accession	<a href="#">O19002</a>
Sequence	<b>Biotin-QGRKRRQ(pT)SMTDFYH</b>

**Biotinylated p21Cip1-T145 Phospho Peptide - Additional Information****Gene ID** 1026**Other Names**

Cyclin-dependent kinase inhibitor 1, CDK-interacting protein 1, Melanoma differentiation-associated protein 6, MDA-6, p21, CDKN1A, CAP20, CDKN1, CIP1, MDA6, PIC1, SDI1, WAF1

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Biotinylated p21Cip1-T145 Phospho Peptide - Protein Information****Name** CDKN1A ([HGNC:1784](#))**Function**

Plays an important role in controlling cell cycle progression and DNA damage-induced G2 arrest (PubMed:<<http://www.uniprot.org/citations/9106657>>9106657</a>). Involved in p53/TP53 mediated inhibition of cellular proliferation in response to DNA damage. Also involved in p53-independent DNA damage-induced G2 arrest mediated by CREB3L1 in astrocytes and osteoblasts (By similarity). Binds to and inhibits cyclin-dependent kinase activity, preventing phosphorylation of critical cyclin-dependent kinase substrates and blocking cell cycle progression. Functions in the nuclear localization and assembly of cyclin D-CDK4 complex and promotes its kinase activity towards RB1. At higher stoichiometric ratios, inhibits the kinase activity of the cyclin D-CDK4 complex. Inhibits DNA synthesis by DNA polymerase delta by competing with POLD3 for PCNA binding (PubMed:<<http://www.uniprot.org/citations/11595739>>11595739</a>).

**Cellular Location**

Cytoplasm. Nucleus

**Tissue Location**

Expressed in all adult tissues, with 5-fold lower levels observed in the brain

**Biotinylated p21Cip1-T145 Phospho Peptide - Images**