

Biotinylated Cdk7-T170 Non-phospho Control Peptide
Synthetic Peptide
Catalog # SP2038d**Specification**

Biotinylated Cdk7-T170 Non-phospho Control Peptide - Product Information

Primary Accession	P50613
Other Accession	Q03147
Sequence	Biotin-GSPNRAYTHQVVTRW

Biotinylated Cdk7-T170 Non-phospho Control Peptide - Additional Information**Gene ID** 1022**Other Names**

Cyclin-dependent kinase 7, 39 kDa protein kinase, p39 Mo15, CDK-activating kinase 1, Cell division protein kinase 7, Serine/threonine-protein kinase 1, TFIIH basal transcription factor complex kinase subunit, CDK7, CAK, CAK1, CDKN7, MO15, STK1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Biotinylated Cdk7-T170 Non-phospho Control Peptide - Protein Information**Name** CDK7**Synonyms** CAK, CAK1, CDKN7, MO15, STK1**Function**

Serine/threonine kinase involved in cell cycle control and in RNA polymerase II-mediated RNA transcription (PubMed: [9852112](http://www.uniprot.org/citations/9852112), PubMed: [19136461](http://www.uniprot.org/citations/19136461), PubMed: [26257281](http://www.uniprot.org/citations/26257281), PubMed: [28768201](http://www.uniprot.org/citations/28768201)). Cyclin-dependent kinases (CDKs) are activated by the binding to a cyclin and mediate the progression through the cell cycle. Each different complex controls a specific transition between 2 subsequent phases in the cell cycle. Required for both activation and complex formation of CDK1/cyclin-B during G2-M transition, and for activation of CDK2/cyclins during G1-S transition (but not complex formation). CDK7 is the catalytic subunit of the CDK-activating kinase (CAK) complex. Phosphorylates SPT5/SUPT5H, SF1/NR5A1, POLR2A, p53/TP53, CDK1, CDK2, CDK4, CDK6 and CDK11B/CDK11 (PubMed: [9852112](#)

[9372954](http://www.uniprot.org/citations/9372954), PubMed: [9840937](http://www.uniprot.org/citations/9840937), PubMed: [19136461](http://www.uniprot.org/citations/19136461), PubMed: [26257281](http://www.uniprot.org/citations/26257281), PubMed: [28768201](http://www.uniprot.org/citations/28768201)). Initiates transcription by RNA polymerase II by mediating phosphorylation of POLR2A at 'Ser-5' of the repetitive C- terminal domain (CTD) when POLR2A is in complex with DNA, promoting dissociation from DNA and initiation (PubMed: [19136461](http://www.uniprot.org/citations/19136461), PubMed: [26257281](http://www.uniprot.org/citations/26257281), PubMed: [28768201](http://www.uniprot.org/citations/28768201)). CAK activates the cyclin-associated kinases CDK1, CDK2, CDK4 and CDK6 by threonine phosphorylation, thus regulating cell cycle progression. CAK complexed to the core-TFIIF basal transcription factor activates RNA polymerase II by serine phosphorylation of the CTD of POLR2A, allowing its escape from the promoter and elongation of the transcripts (PubMed: [9852112](http://www.uniprot.org/citations/9852112)). Its expression and activity are constant throughout the cell cycle. Upon DNA damage, triggers p53/TP53 activation by phosphorylation, but is inactivated in turn by p53/TP53; this feedback loop may lead to an arrest of the cell cycle and of the transcription, helping in cell recovery, or to apoptosis. Required for DNA-bound peptides-mediated transcription and cellular growth inhibition.

Cellular Location

Nucleus. Cytoplasm. Cytoplasm, perinuclear region. Note=Colocalizes with PRKCI in the cytoplasm and nucleus (PubMed:15695176). Translocates from the nucleus to cytoplasm and perinuclear region in response to DNA-bound peptides (PubMed:19071173).

Tissue Location

Ubiquitous.

Biotinylated Cdk7-T170 Non-phospho Control Peptide - Images