

**Interleukin - 6 Receptor Peptide**  
**Synthetic Peptide**  
**Catalog # SP2442a****Specification**

---

**Interleukin - 6 Receptor Peptide - Product Information**Primary Accession  
Sequence[P08887](#)  
NH2-TSLPVQDSSSVP-COOH**Interleukin - 6 Receptor Peptide - Additional Information****Gene ID** 3570**Other Names**

Interleukin-6 receptor subunit alpha, IL-6 receptor subunit alpha, IL-6R subunit alpha, IL-6R-alpha, IL-6RA, IL-6R 1, Membrane glycoprotein 80, gp80, CD126, IL6R

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Interleukin - 6 Receptor Peptide - Protein Information****Name** IL6R ([HGNC:6019](#))**Function**

Part of the receptor for interleukin 6. Binds to IL6 with low affinity, but does not transduce a signal (PubMed:<a href="http://www.uniprot.org/citations/28265003" target="\_blank">28265003</a>). Signal activation necessitate an association with IL6ST. Activation leads to the regulation of the immune response, acute-phase reactions and hematopoiesis (PubMed:<a href="http://www.uniprot.org/citations/30995492" target="\_blank">30995492</a>, PubMed:<a href="http://www.uniprot.org/citations/31235509" target="\_blank">31235509</a>). The interaction with membrane-bound IL6R and IL6ST stimulates 'classic signaling', the restricted expression of the IL6R limits classic IL6 signaling to only a few tissues such as the liver and some cells of the immune system. Whereas the binding of IL6 and soluble IL6R to IL6ST stimulates 'trans- signaling'. Alternatively, 'cluster signaling' occurs when membrane- bound IL6:IL6R complexes on transmitter cells activate IL6ST receptors on neighboring receiver cells (Probable).

**Cellular Location**

[Isoform 1]: Cell membrane {ECO:0000250|UniProtKB:P22272}; Single-pass type I membrane protein [Soluble interleukin-6 receptor subunit alpha]: Secreted

**Tissue Location**

[Isoform 2]: Expressed in peripheral blood mononuclear cells and weakly found in urine and serum. 1%-20% of the total sIL6R in plasma is generated by alternative splicing (PubMed:28060820).

**Interleukin - 6 Receptor Peptide - Images**