

[Asn370] tyrosinase (368 - 376) Synthetic Peptide Catalog # SP2477b

Specification

[Asn370] tyrosinase (368 - 376) - Product Information

Primary Accession Other Accession Sequence <u>Q9BDE0</u> <u>P14679, Q8MIU0, P54834</u> NH2-YMNGTMSQV-COOH

[Asn370] tyrosinase (368 - 376) - Additional Information

Other Names Tyrosinase, Monophenol monooxygenase, TYR

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

[Asn370] tyrosinase (368 - 376) - Protein Information

Name TYR

Function

This is a copper-containing oxidase that functions in the formation of pigments such as melanins and other polyphenolic compounds (By similarity). Catalyzes the initial and rate limiting step in the cascade of reactions leading to melanin production from tyrosine (By similarity). In addition to hydroxylating tyrosine to DOPA (3,4- dihydroxyphenylalanine), also catalyzes the oxidation of DOPA to DOPA- quinone, and possibly the oxidation of DHI (5,6-dihydroxyindole) to indole-5,6 quinone (By similarity).

Cellular Location

Melanosome membrane {ECO:0000250|UniProtKB:P14679}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P14679}. Melanosome {ECO:0000250|UniProtKB:P11344}. Note=Proper trafficking to melanosome is regulated by SGSM2, ANKRD27, RAB9A, RAB32 and RAB38 {ECO:0000250|UniProtKB:P11344}

[Asn370] tyrosinase (368 - 376) - Images